opendaccess

# **Evaluation of Knowledge Level of Under Five Children's Mother Regarding Vitamin A Deficiency**

Daneshwari Hiremath<sup>1</sup>, Deelip S Natekar<sup>2</sup>, Mahalaxmi C<sup>3</sup>, Kalakappa N Huli<sup>3</sup>, Meenakshi M<sup>3</sup>\*, Manojkumar CH<sup>3</sup>, Roopa J<sup>3</sup>, Apoorva A<sup>3</sup>

<sup>1</sup>Associate Professor and HOD, Department of Child Health Nursing, Shri B.V.V.S. Sajjalashree Institute of Nursing Sciences, Navangar, Bagalkot, Karnataka, India

<sup>2</sup>Principal, Department of Community Health Nursing, Shri B.V.V.S. Sajjalashree Institute of Nursing Sciences, Navangar, Bagalkot, Karnataka, India

<sup>3</sup>Student, Department of Child Health Nursing, Shri B.V.V.S. Sajjalashree Institute of Nursing Sciences, Navangar, Bagalkot, Karnataka, India

\*Address for Correspondence: Meenakshi M, Student, Department of Child Health Nursing, Shri B. V. V. S. Sajjalashree Institute of Nursing Sciences, Navanagar, Bagalkot- 587102, Karnataka, India E-mail: manjulasc9686@gmail.com

# Received: 26 Nov 2023/ Revised: 23 Dec 2023/ Accepted: 03 Feb 2024

## ABSTRACT

Background: Early childhood is regarded as one of the most essential and valuable periods before attaining adulthood. The first five years of life are the most important for a child's development as many biological, psychological, and emotional changes take place during this time. Children need enough nutrients during the developmental stage to support growth and immunity. They experience certain deficiency disorders if their needs are not fulfilled. Therefore, the current study was undertaken to establish a theoretical basis for the assessment of knowledge of mothers of under five years regarding vitamin A deficiency.

Methods: The study utilized a descriptive survey design and a simple random technique was used to obtain 60 samples from Anganawadi, urban area of Bagalkot, Karnataka, India. Data were collected using a semi-structured questionnaire to assess the level of knowledge among mothers of under-five children regarding vitamin A deficiency. The collected data were analysed by using descriptive and inferential statistics.

Results: Among the 60 samples studied, the knowledge of under-five children's mothers was very poor for 10% of mothers, poor for 66.66% of mothers, and good knowledge was reported for 23.33% mothers. The results of the present study showed a nonsignificant association between knowledge levels of mothers of under-five children and age group (p<1), religion (p=0), education (p<0.139), number of children (p<0.262), income (p<0.258), occupation (p<0.686), and type of family (p<0.022).

Conclusion: The study concluded that the knowledge of mothers of under-five children was very poor for 10% of mothers and poor for 66.66% of mothers.

Key-words: Children, Childhood blindness, Maternal Awareness, Vitamin A deficiency, Knowledge, Mothers

## INTRODUCTION

Vitamin A is critical for the optical system's normal functioning. It is also required to maintain cell function for growth, production of red blood cells, epithelial cells,

How to cite this article

Hiremath D, Natekar DS, Mahalaxmi C, Huli KN, Meenakshi M, et al. Evaluation of Knowledge Level of Under Five Children's Mother Regarding Vitamin A Deficiency. SSR Inst Int J Life Sci., 2024; 10(2): 5174-5179



Access this article online https://iijls.com/

reproduction, and immunity. Vitamin A deficiency is an essential factor for childhood blindness and a significant contributing factor to mortality and morbidity among under-five children <sup>[1]</sup>.

It is mainly seen in poorer people and under developing countries. Two key factors leading to vitamin A deficiency in children are: 1) Their mothers are deficient in vitamin A, which causes the production of breast milk low in vitamin A; and 2) Children are weaned onto Vitamin A- deficient diet <sup>[2]</sup>. Their prolonged illness during childhood, which is exacerbated by anorexia,

**Research Article** 

malabsorption, and accelerated catabolism, is a third factor that contributes to their low vitamin A level <sup>[3,4]</sup>.

A mother must know the causes, importance, sources, and deficiency of vitamins. According to our study, 10% of mothers of under-five children have very poor knowledge, 66.66% of mothers have poor knowledge, and 23.33% mothers have good knowledge regarding vitamin A deficiency. Vitamin A deficiency is a systemic condition that mostly affects the eyes. It arises from diets not providing enough vitamin A to support growth and development, psychological functions, and periods of added stress due to illness <sup>[5]</sup>. Vitamin A is found in foods such as milk, red and orange fruits, liver, red palm oil, eggs, and green leafy vegetables. Although these sources provide varying amounts of vitamin A to the body <sup>[6]</sup>.

### MATERIALS AND METHODS

A descriptive survey design was utilized for the present study. A simple random technique obtained 60 samples from the Anganawadi urban area of Bagalkot, Karnataka, India. This study was conducted in selected urban areas of Bagalkot. The study setting was selected according to availability of mothers' the and investigators' convenience. Data were collected from mothers of under-five children using semi-structured а questionnaire and their knowledge regarding vitamin A deficiency was evaluated. Descriptive and inferential statistics were used to analyse the collected data.

Research Design- A descriptive survey design was used to assess data from under-five children's mothers about vitamin A deficiency. The dependent variable was used to determine the level of knowledge about Vitamin A deficiency in mothers of children under five years old. In contrast, the independent variable was used to assess the mothers' understanding of preventing vitamin A children under five. deficiency in Various sociodemographic variables of mothers of children of 0-5 years of age, including mother's age, educational status, religion, monthly income, total number of children, occupation, and type of family, were studied. The present study sample size was 60 mothers of children under five who resided in urban areas of Bagalkot, Karnataka, India. A convenient sampling technique selected the samples. The mothers were chosen conveniently according to the duration of study. Those

subjects who fulfilled the inclusion criteria were selected using a simple random technique. A semi-structured questionnaire was utilised to collect the data in the present study.

A pilot study was carried out at the end of the planning phase to explore and test the research elements. A pilot study was conducted in some selected urban area of Bagalkot from 28-6-2023 to 29-6-2023 to determine the study design's feasibility and practicability. Six mothers were chosen randomly using self-administered structured closed-ended knowledge and practice questionnaires.

**Statistical Analysis-** The obtained data were assessed using inferential and descriptive statistics. Sociodemographic data were evaluated by using percentage and frequency distribution. Mean and standard deviation were used to analyse the scores of under-five children's mothers. The chi-square test was used to assess the relationship between the knowledge of under-five children's mothers and the selected sociodemographic variables.

**Ethical Consideration**- Formal permission was taken from the Principal, BVVS Sajjalashree Institute of Nursing Sciences, Bagalkot, Karnataka, India. An ethical clearance certificate was obtained and enclosed from the ethical committee of B.V.V.S Sajjalashree Institute of Nursing Sciences, Bagalkot, Karnataka, India. Researchers explained the purpose of the study to the participants, and written informed consent was obtained from the mothers of the children under five participating in the study. Anonymity and confidentiality regarding the data and identity of mothers were maintained.

### RESULTS

The results of the present study show that among 60 samples, the knowledge level of under-five children's mothers was very poor for 10% mothers, poor for 66.66% mothers, and good for 23.33% mothers. The frequency and percentage distribution of sociodemographic characteristics of mothers of under-five children is shown in Table 1. The majority (53.33%) of mothers were 24-29 years old, Hindu (93.33%), housewives (96.66%), living in a joint family (51.66%), had 2 children (55%), with a monthly income of Rs.5000-10000 (83.33%).

Crossef DOI: 10.21276/SSR-IIJLS.2024.10.2.24

Sociodemographic variables	Frequency	Percentage (%)				
Age of mother (in years)						
18-23	9	15				
24-29	32	53.33				
30-35	17	28.33				
36 and above	2	3.33				
Religion of mother						
Hindu	56	93.33				
Muslim	4	6.66				
Christian	0	0				
Other	0	0				
Educational status of the mother						
Primary	24	40				
PUC	23	38.33				
Degree	9	15				
No formal education	4	6.66				
	Total number of children					
1 child	12	20				
2 children	33	55				
3 children	13	21.66				
More than 3 children	2	3.33				
Fami	ly monthly income (in Rupees)					
5000–10000	50	83.33				
10000–15000	9	15				
15000–20000	1	1.66				
20000 and above	0	0				
	Occupation of mother					
Housewife	58	96.66				
Business	1	1.66				
Government servant	0	0				
Private employees	1	1.66				
Type of family						
Joint	31	51.66				
Nuclear	29	48.33				

# **Table 1**: Percentage distribution and frequency of sociodemographic variables

As shown in Table 2, the calculated mean was 10.466, the standard error was 0.0513, and the standard difference was 0.394; hence, the knowledge of underfive children's mothers was insignificant. Percentagewise distribution scale of the level of knowledge of children's mothers is given in Table 3. Out of 60 samples, six (10%) have very poor knowledge, 40 (66.66%) have poor knowledge, and 14 (23.33%) have good knowledge.

crossef DOI: 10.21276/SSR-IIJLS.2024.10.2.24

Table 2. Knowledge about vitamin A denderely in mothers of duder five emidren					
Test	Mean	Standard error	Standard difference	Table value	
	10.466	0.0513	0.394	3.84	

 Table 2: Knowledge about vitamin-A deficiency in mothers of under-five children

Table 3: Scale of the level of knowledge of under-five children's mothers

Range of score	Knowledge level	Frequency	Percentage (%)
0-6	Very poor	6	10
7-12	Poor	40	66.66
13-18	Good	14	23.33
19-24	Very good	0	0
25-30	Excellent	0	0

There was no significant association between the age group of under-five children's mothers and their knowledge about vitamin A deficiency because the calculated value (1) was less than the table value (3.846). Non-significant association was established between the religion of all mothers of under-five children and their knowledge level because the study value (0) was less than the table value (3.846). Similarly, a non-significant relationship was observed between the educational status of mothers (study value=0.1395), number of children (study value=0.2622), family income (study value=0.2586), occupation of mothers (study value=0), type of family (study value=0.022) and their knowledge because study value for each variable was less than table value (3.846) (Table 4).

Socio-demographic variables	Chi-square calculated value	Association
Age of mother	1	NS
Religion	0	NS
Educational status	0.1395	NS
Number of children	0.2622	NS
Income	0.2586	NS
Occupation	0	NS
Type of family	0.022	NS

Table 4: Relationship between knowledge of mothers and the selected sociodemographic variables

Chi-Square table value=3.846; Degree of freedom (DF)=1; NS: non-significant.

## DISCUSSION

The present study evaluated the knowledge levels of under-five children's mothers regarding vitamin A deficiency in selected urban areas of Bagalkot, Karnataka, India. It was observed that many mothers (53.33%) belong to the age group of 24-29 years, while only two (3.33%) mothers belong to the 36 and above age group. In comparison with a similar study, it was found that many mothers (40%) belong to the 21-25 years age group and only 8 (13.3%) mothers belong to the below 20 years age group <sup>[7]</sup>.

Most mothers (93.33%) in the present study were Hindu by religion, while no one (0%) was Christian. Comparable results were obtained by Castro *et al.*, who found 6% of mothers from the Hindu religion and 18% from the Muslim community <sup>[8]</sup>. The educational status of most mothers in the present study was primary (40%) and only 6.66% mothers obtained nonformal education. On comparing the data to previous studies, Mesh *et al.* found that 22 (36.70%) mothers belonged to primary education and 8 (13.30%) mothers were postgraduate <sup>[9]</sup>. Many mothers (55%) have 2 children, while only 2 (3.33%) have more than 3 children. Similar results were reported by Miglioli et al.; they found that 27(45%) mothers have 2 children, while 7(17.70%) mothers have more than 3 children <sup>[10]</sup>.

The monthly family income of most mothers (83.33%) in the present study was Rs.5000-10000 while none of the mother's family income was more than Rs.20000 and above. Similar results were reported by a previous study where 21(35%) mothers' family income was below Rs.20000 and 8(13.30%) mothers had a family monthly income of more than Rs.20000 [11]. Many mothers were housewives, 58(96.66%) and none were government employees (0%). Haslow et al. [12] in a similar study determined that 38(63.30%) mothers were housewives and 7(12.40%) were private employees. In the present study, most mothers (51.66%) belong to a joint family, and 29(48.33%) belong to nuclear research. These findings were contrary to those of a study by Arlappa <sup>[13]</sup> who found that 5(8%) of mothers belong to a joint family and 36 (92%) of mothers belong to a nuclear family. Few studies have also assessed vitamin A dietary intake among children under five years old to evaluate vitamin A supplementation coverage. One such study by Yadav et al. conducted among children of 6-60 months in Anganwadis of Chandigarh revealed that only 20% children took food and vegetables rich in vitamin A during the past 24 hours. We did not consider this dimension in our study. However, such a targeted approach is required to update the current nutrition programme [14].

Out of the total samples studied, 10% have very poor knowledge, 66.66% have poor knowledge, and 23.33% have good knowledge. Another study by Yamunambigai *et al.* revealed that out of 50 mothers of under-five children, the majority (72%) had insufficient knowledge. In contrast, 16% of them had fair knowledge and 12% of mothers had adequate knowledge regarding vitamin A deficiency <sup>[15]</sup>. Few studies reported similar findings, where most mothers with under-five children had an average poor knowledge of vitamin A deficiency <sup>[16,17]</sup>.

Policy changes are currently required in India regarding the national programme to choose targeted distribution over universal distribution. In addition to health measures, the National Institute for Transforming India suggests bolstering the national programme for control of vitamin A deficiency through six months of vitamin A supplementing in addition to medical intervention <sup>[18]</sup>. Apart from short-term emergency measures of administration of vitamin A to the vulnerable group, a long-term solution through home gardening and nutrition education programmes can be initiated, as opined by DeMaeyer<sup>[19]</sup>.

## CONCLUSIONS

Children are a vulnerable segment in terms of vitamin A deficiency. The present study concluded that mothers of children under five had inadequate knowledge of vitamin A deficiency. Also, there was a non-significant relationship between the studied sociodemographic variables and the level of knowledge of mothers of children under five.

Under-five children are more prone to vitamin A deficiency diseases such as blindness, Crohn's disease, xeropthalmia, and visual impairment. Giving proper knowledge and education to the mothers about the importance of vitamin A will reduce the risk of vitamin A deficiency in under-five children. The present study will be useful to build awareness among mothers regarding vitamin A deficiency.

# ACKNOWLEDGMENTS

We thank the anonymous referees for their useful suggestions. Words are insufficient to express my heartfelt thanks and sincere gratitude towards those helping hands.

### **CONTRIBUTION OF AUTHORS**

Research concept- Dr Dileep Natekar, Dr. Daneshwari Hiremath Research design- Dr. Daneshwari Hiremath Supervision- Dr. Daneshwari Hiremath Materials- All Researchers Data collection- All Researchers Data analysis and interpretation- All Researchers Literature search- All Researchers Writing article- All researchers Writing article- All researchers Critical review-. Dr. Daneshwari Hiremath Article editing-. Dr. Daneshwari Hiremath Final approval- Dr. Daneshwari Hiremath

#### REFERENCES

 Rozario MD, Costa ND, Akter H. Interpersonal interaction and hindrances of nursing students in clinical learning. Asian Journal of Nursing Education and Research, 2023; 13(3): 171-78.

- [2] Varghese S, Manuel S, Tessya A. A study to assess the knowledge of mothers of underfive children regarding the importance of Vitamin A among selected areas of Pallithottam, Kollam. Asian J., 2020; 10(1): 84-88.
- [3] Sindhu P, Nirmala R. A study on Vitamin A deficiency. Int J Res., 2020; 7(10): 101-10.
- [4] Ayoya MA, Bendech MA, Baker SK, Ouattara F, et al. Determinants of High Vitamin A Supplementation Coverage among Pre-School Children in Mali: The National Nutrition Weeks Experience. J Public Health Nutr., 2007; 10(11): 1241-46.
- [5] Sangho H, Belem B, Keita HD, Keita AS. Vitamin A supplementation in children under five during a oneweek nutrition intensification program in Mali. Santee Publique., 2013; 25(6): 821-27.
- [6] Njue MW, Makokha AO, Mutai JK. Vitamin A supplementation awareness among mothers of children under five years old at Mbagathi district hospital, Nairobi, Kenya. East Afr J Public Health., 2010; 7(3): 233-41.
- [7] Kamau MW, Makokha AO, Mutai JK, Mugoya Ik. Factors influencing Vitamin A supplementation among mothers of children under five years old at Mbagathi District Hospital, Kenya. East Afr Med J., 2012; 89(4): 134-41.
- [8] Castro IRR, Normand P, Ferias DR, Berti TL, Schinhaglia RM, et al. Factors associated with anaemia and Vitamin A deficiency in Brazilian children under 5 years old: Brazilian National Survey On Child Nutrition (ENANI-2019). Cad Saude Publicantion, 2023; 39(2): 309-14.
- [9] Mesh Rams II, Née Raja G, Longvah T. Vitamin A deficiency, anemia, and nutritional status of under 5year children from Northeast India. Indian J Community Med., 2021; 46(4): 673-79.
- [10] Miglioli TC, Fonseca VM, Gomes Junior SC, de Lira PIC, Filho MB. Vitamin A deficiency in mothers and children in the State of Pernambuco. Cien Saudi Colet., 2013; 18(5): 1427-40.

- [11]Ahenda P, Washburn DJ, Colwell B, Lee S, Gwarzo I, et al. Shifts in social determinants of Vitamin A supplementation among children under five in Kenya. Matern Child Health J., 2023; 27(7): 1284-92.
- [12] Haselow NJ, Joshi V, Bayo PN, Murye JW, Shaban SN, et al. A review of Vitamin A supplementation in South Sudan: Successes, challenges, and opportunities for the way forward. Glob Health Sci Pract., 2022; 10(3): 21-30.
- [13]Arlappa N. Vitamin A deficiency is still a public health problem in India. Indian J Pediatr., 2011; 48(: 853-54.
- [14]Yamunambigai M, Manohari K, Nithiya K. A descriptive study to assess the knowledge regarding vitamin A deficiency among mothers of under-five children in selected rural areas at Puducherry. International Journal of Advanced Research, 2023; 11: 685-91.
- [15]Bhatia A. A descriptive study to assess the knowledge regarding Vitamin A deficiency disorders among the mothers of underfive children in selected rural areas of district Ludhiana, Punjab. Int J Nur Edu Res., 2017; 5(4): 395-98.
- [16] Vijayaraghavan K. National Control Programme against nutritional blindness due to Vitamin A deficiency. Indian J Med Res., 2018; 148(5): 496-502.
- [17]Demaeyer EM. The WHO Programme of prevention and control of Vitamin A deficiency, exophthalmia and nutritional blindness. Nutr Health., 1986; 4(2): 105-12.
- [18]Soares-Mota M, Silva TA, Gomes LM, Pinto MA, Mendonca LM, et al. High prevalence of Vitamin A deficiency in Cohn's disease patients according to serum retinol levels and the relative dose-response test. World J Gastroenterol., 2015; 21(5): 1614-20.
- [19]Yadav R, Patial A, Bharti B, Attri SV, Bhatia P. Dietary Vitamin A intake, coverage of Vitamin A megadose supplementation, and prevalence of Vitamin A deficiency among marginalized children 6-59 months in Anganwadis of Chandigarh: A multistage cluster sampling survey. Indian J Community Med., 2021; 46(4): 692-96.

**Open Access Policy:** 

Authors/Contributors are responsible for originality, contents, correct references, and ethical issues. SSR-IIJLS publishes all articles under Creative Commons Attribution- Non-Commercial 4.0 International License (CC BY-NC). <u>https://creativecommons.org/licenses/by-nc/4.0/legalcode</u>