

# Exploring the Wound Healing Potential of *Tridax procumbens* Extract: A Comprehensive Analysis through *in vitro* Cytotoxicity and Scratch Assay Studies

Nidhi Tripathi<sup>1\*</sup>, Divyanshi Makode<sup>2</sup>, Vishvam Jawade<sup>2</sup>, Mahi Verma<sup>2</sup>, Jagat Rakesh Kanwar<sup>3</sup>, Rashmi Chowdhary<sup>4</sup>, Anjali Choudhary<sup>5</sup>, Preeti Chandurkar<sup>6</sup>, Ashish Kumar Yadav<sup>7</sup>, Vrushali D. Patil<sup>7</sup>

<sup>1</sup>Mentor, E YUVA Team, E YUVA Centre, Career College (Autonomous), Bhopal, India

<sup>2</sup>E YUVA Fellow, E YUVA Centre, Career College (Autonomous), Bhopal, India

<sup>3</sup>Professor & Head, Dept. of Biochemistry, AIIMS, Bhopal, India

<sup>4</sup>Professor, Dept. of Biochemistry, AIIMS, Bhopal, India

<sup>5</sup>HOD, Dept. of Biotechnology, Career College (Autonomous), Bhopal, India

<sup>6</sup>Project Coordinator, E YUVA Centre, Career College (Autonomous), Bhopal, India

<sup>7</sup>Research Scholar, Dept. of Biochemistry, AIIMS, Bhopal, India

**\*Address for Correspondence:** Dr. Nidhi Tripathi, Mentor, E YUVA Team, E YUVA Centre, Career College (Autonomous), Bhopal, Madhya Pradesh, India

**E-mail:** [nidhibiochemistry@gmail.com](mailto:nidhibiochemistry@gmail.com)

Received: 11 Nov 2023 / Revised: 12 Dec 2023 / Accepted: 13 Feb 2024

## ABSTRACT

**Background:** Wound healing, a complex biological process, is crucial for preventing infections and maintaining skin integrity. This comprehensive review examines wound dynamics based on location, etiology, symptoms, and clinical appearance. Emphasizing the importance of precise progression through inflammation, cellular proliferation, and remodeling phases, the study highlights the risk of chronic wounds with delays or interruptions.

**Methods:** This study explores natural alternatives to synthetic wound healing drugs, focusing on *T. procumbens*. Scientific investigations reveal its diverse pharmacological activities, particularly in ointment formulations. *In vitro* cytotoxicity and scratch tests, using the MTT test on Vero-E6 cell lines, demonstrate *T. procumbens*' potential for wound healing. The research aims to address the challenges posed by the side effects of synthetic drugs.

**Results:** The research provides insights into skin tissue regeneration through *T. procumbens*, revealing its ability for wound healing. The cytotoxicity index is calculated after applying drug concentrations, and regression analysis yields the IC<sub>50</sub>, offering information on the effect on cell viability.

**Conclusion:** Combining traditional wisdom with contemporary research, this study bridges the gap between technology and tradition, presenting a comprehensive method for wound management. *T. procumbens* emerges as a promising herbal treatment, with its potential demonstrated through diverse pharmacological activities and cellular reactions.

**Key-words:** Cellular proliferation, Chronic wounds, Cytotoxicity, Inflammation, Medicinal plants, Remodeling phases, *Tridax procumbens*, Wound healing

## How to cite this article

Tripathi N, Makode D, Jawade V, Verma M, Kanwar JR, et al. Exploring the Wound Healing Potential of *Tridax procumbens* Extract: A Comprehensive Analysis through *in vitro* Cytotoxicity and Scratch Assay Studies. SSR Inst Int J Life Sci., 2024; 10(2): 5146-5151.



Access this article online

<https://ijls.com/>

## INTRODUCTION

Wound healing is a sophisticated process involving tissue restoration following injury or damage. A wound is described as disrupting the normal continuity of the skin, leading to a break in its cellular and anatomical structures, thereby affecting its regular functionality. This intricate process can be classified based on the wound's location, origin, presenting symptoms, type of injury, depth, tissue loss, or clinical appearance. A comprehensive understanding of the multifaceted

nature of wounds is crucial for developing effective strategies for their management and promoting optimal healing outcomes.

The healing process can be conceptualized into three overlapping phases: inflammation, cellular proliferation, and remodelling. These phases are carefully orchestrated to ensure the restoration of normal tissue function <sup>[1]</sup>. During the initial inflammatory phase, the body responds to the injury by eliminating pathogens and cellular debris through the influx of cytokines and growth factors. Subsequently, the cellular proliferation phase involves the formation of granulation tissue through processes like angiogenesis, with keratinocytes and fibroblasts migrating to the wound site. The maturation phase is marked by restoring the skin barrier, repairing granulation tissue within the scar, and vessel regression <sup>[2]</sup>. This newly formed tissue gradually gains tensile strength, with collagen production, cross-linking, and reorganization occurring over an extended period <sup>[3]</sup>. The accurate and regular progression of these phases is crucial, as interruptions or prolongations can lead to chronic wounds or delayed healing, often associated with conditions such as ischemia, diabetes, and hypertension <sup>[4]</sup>.

A wound disrupts cellular, anatomical, and functional continuity in living tissue resulting from various insults, including physical, chemical, thermal, microbial, or immunological insults. Essentially, a wound is a break in epithelial integrity accompanied by disruption in the structure and function of underlying normal tissue <sup>[5-8]</sup>. While the natural healing process of wounds initiates and progresses, the outcomes are not always favorable <sup>[9]</sup>.

To enhance the quality and speed of the healing process, synthetic drugs like nitrofurazone, gentamicin, and mupirocin are commonly used, which have many side effects. In recent years, there has been a growing trend towards using natural medicines to mitigate these drawbacks <sup>[10]</sup>. Many medicinal plants, with a long history of efficacy in various diseases, have been explored for their potential in wound healing. However, despite their traditional use, there is often a lack of scientific data or information to prove their effectiveness, understand their active ingredients, or elucidate their mechanisms of action <sup>[11]</sup>.

In this context, *T. procumbens* leaves emerge as one of the medicinal plants traditionally used for wound

healing, particularly by tribal communities. *T. procumbens*, commonly known as "Coat buttons" or "Mexican daisy," is a weed plant belonging to the Asteraceae family. It is often found in waste places, roadsides, and hedges. The leaves of *T. procumbens* contain various components such as 26% crude protein, 17% crude fibre, 39% soluble carbohydrates, 5% calcium oxide, and 5% Luteolin, glucoluteolin, and quercetin. Traditionally, the matured leaves are crushed to make a paste and applied to the surface of wounds. Additionally, these leaves have been used for various ailments like high blood pressure, bronchial catarrh, dysentery, malaria, diarrhoea, and to prevent hair fall <sup>[12-14]</sup>. Scientific investigations into *T. procumbens* have revealed a wide range of pharmacological activities, including antioxidant, antimicrobial, anti-inflammatory, vasorelaxant, anti-leishmanial, anti-arthritis, antihypertensive, hepatoprotective, anticancer, hemostatic, anti-anemic, and anti-diabetic properties <sup>[15]</sup>. Cell culture techniques are indispensable in developing biotechnology products, as they provide quick setups and a wide range of in vitro tests such as proliferation, migration, viability, inflammation, and apoptosis <sup>[11]</sup>. The comparison of cell migration outcomes using *different in vitro* wound healing techniques provides valuable insights into the intricate interplay between cellular responses and external factors <sup>[16]</sup>. The present study aims to harness the potential of *T. procumbens* extract for wound healing.

## MATERIALS AND METHODS

**Formulation of extract-** *T. procumbens* plants were gathered from Misrod Bhopal farms to formulate the gel. To prepare the extract, the flowers from the collected plants were removed, and the remaining components were cleaned with tap water, distilled water, and filter paper under the sun for seven days. Following the entire drying of the plants, the powdered leaves, stems, and roots were gathered and put through sieve No. 12 to eliminate any remaining debris. Following this, a 1:10 ratio of powder to water was used for dissolution. This mixture was incubated for a whole day. After that, it was filtered, and the filtrate was saved for the water to evaporate <sup>[17]</sup>.

**Cytotoxicity Test-** To assess the potential toxicity of *T. procumbens* extract on cells. In brief, 5000 cells per well

were seeded in 96-well plates, and the cells were exposed to the ointment for 24-72 hours, with each condition performed in triplicate. Cell growth was evaluated using MTT dye. 100  $\mu$ L of MTT dye (0.5 mg working stock) was added to each well and incubated for 4 hours. Following incubation, the formazan crystals formed within the cells were solubilized using dimethyl sulfoxide, and the optical density was measured at 570 nm using a spectrophotometer <sup>[18]</sup>. The percentage of cell growth cells was determined using the following formula:

## RESULTS










The drug-containing *T. procumbens* extract was dissolved in Acetone. The IC<sub>50</sub> was calculated using MTT assay. The range of IC<sub>50</sub> was 110 $\mu$ g to 125.5  $\mu$ g. Accordingly, two different concentrations of the drug were used 4 $\mu$ g/ $\mu$ L & 10 $\mu$ g/ $\mu$ L. 10 $\mu$ L of both concentrations were added to 24-

$$\text{Cell growth (\%)} = \frac{\text{A570/630 of treated cells}}{\text{A570/630 of control cells}}$$


**Scratch assay-** To determine the effect of the compound on cell growth, scratch assay was performed. In the present study, 5000 cells/well were seeded in 96-well plates, and cells were treated with *T. procumbens* extract for 24-72 hours (in triplicate). Cell growth was measured using MTT dye and incubated for 4 hours. The percentage of cell growth cells will be determined using the formula: cell growth (%) = A570/630 of treated cells/A570/630 of control cells <sup>[18]</sup>.

well plate. The final drug concentration in each well was 40 $\mu$ g & 100  $\mu$ g respectively. The result of the scratch assay depicted the effectiveness of both concentrations. But, the 40 $\mu$ g drug concentration was found to be more effective (Table 1).

**Table 1:** Cell migration appeared between scratch area in Vero E 6 Cell

Intervals	Concentration of Ointment (Vero E6 cell line)		Control Vero E6 cell line
	40 ug (10ug/ $\mu$ L)	100 ug (10ug/ $\mu$ L)	
Day 0			
Day 1			
Day 2			





Day 0: Scratch was introduced in vero cell line and after that *T. procumbens* extract was inoculated in the cell lines,  
 Day1: Cell synthesis was started, Day 2: Cell were come closer Day 3: Wound was completely closed.

## DISCUSSION

In the wound healing process, cell migration plays a crucial role in promoting wound closure. The quantification of wound closure following various treatments is illustrated in Table 1. In the present study, vero cells exhibited 99.9% migration toward the scratched wound after 24 hours. It is crucial to note that cell migration is a pivotal event in wound healing, triggering subsequent stages of the curing process. Therefore, investigating vero cells migration after exposure to drug formulations is essential for developing new therapies in wound healing. The scratch assay, a well-established in vitro method for measuring cell migration, assessed the wound-healing efficacy of *T. procumbens* leaf extract.

Numerous studies have been conducted on wound care, emphasizing cutting-edge therapeutic strategies and the creation of plant-based products for treating various wound types. In the excision wound model, Benito *et al.* [19] conducted a study wherein animals were categorized into two groups, each consisting of six animals. The groups were as follows: Group-I (Control), Group-II (Nimesulide), and Group-III (Ethanol extract of *T. procumbens*). Their investigation revealed that administering *T. procumbens* extract, even at doses up to 2000 mg/kg body weight, did not elicit any toxic symptoms or mortality.

No changes in behavioral patterns or indications of toxicity were noted during the 24-hour observation period. As a result, it can be inferred that the extract is safe for subsequent pharmacological evaluations.

The wound scratch assay demonstrated a cell migration rate of approximately  $90 \pm 2\%$  during a 72-hour incubation period, in contrast to the control group, which showed a rate of  $62 \pm 2\%$ . This outcome indicates a notable enhancement in cell migration facilitated by the

herb ointment, leading to the complete disappearance of the scratch area within three days. This accelerated cell migration suggests a potential stimulation of epithelization, proliferation, and collagen viability near the wound region [20].

The ointment formulated with an ethanolic extract of *T. procumbens* demonstrates noteworthy wound healing efficacy, particularly in facilitating wound contraction, especially at a 100 mg/g concentration. However, when compared at the same concentration, standard betadine exhibits higher wound contraction than the test ointment [21].

The ability of *T. procumbens* to cure wounds has been examined in this study. It was found that the *T. procumbens* extract enhanced wound closure in Vero cell lines. Moreover, it was discovered that the extract exhibited no cytotoxic properties. The results of this study suggest that *T. procumbens* could heal wounds and be a good place to locate natural compounds that can do the same. Talekar *et al.* [22] reported that, at a concentration of 20 mg/ml formulation, a notable increase in tertiary and quaternary vessels was observed, attributed to the angiogenic potential of the formulation. The formulation at concentrations of 3 mg/ml and 5 mg/ml demonstrated significant mobilization of keratinocytes and fibroblasts at the injury site. The findings of our current study align with the research conducted by Venkatachalam and Palaniswamy [23]. They observed that the extract of *T. procumbens* leaf demonstrated significant potential as a wound healing agent. Their results indicated a notable reduction in the scratch open area, leading to nearly complete closure of scratches (94%) within 24 hours. Importantly, this effect was achieved without inducing cytotoxicity in L929 mouse fibroblast cells. The difference between the present investigation and Venkatachalam and

Palaniswamy <sup>[19]</sup> was that they studied L929 mouse fibroblasts and in the present study, excellent wound healing properties in *vero* cells was observed. Previous studies reported that the polyherbal formulation exhibited prompt skin regeneration and wound contraction.

Regarding biochemical parameters, there was an elevation in hydroxyproline, hexosamine, and collagen turnover in animals treated with the test drug compared to untreated animals <sup>[22]</sup>. In subsequent studies, it is imperative to conduct further tests adhering to ethical guidelines. These investigations aim to identify optimal treatments for healing wounds and bed sores in diabetic patients.

## CONCLUSIONS

The study highlighting *T. procumbens's* efficacy in promoting wound closure and its non-toxic nature offers a promising avenue for advanced wound care, especially for diabetic patients. Formulating an ointment from its ethanolic extract could significantly benefit diabetic wound management by harnessing the plant's healing properties. Future steps include conducting specific clinical trials on diabetic wounds to confirm efficacy and safety, elucidating the mechanisms through which the plant aids wound healing, and optimizing the formulation for consistent potency. Regulatory approval will be crucial for market introduction, and educating healthcare professionals and patients about this novel treatment option. Additionally, exploring the potential for synergistic effects with other wound-healing agents could lead to even more effective treatments. This approach not only aims to improve healing outcomes for diabetic wounds but also opens up new possibilities for natural, plant-based wound care solutions, potentially transforming the landscape of wound management for diabetic patients and beyond.

## ACKNOWLEDGMENTS

The authors express their sincere appreciation to E-Yuva Centre, Career College (Autonomous) Bhopal, and the Biotechnology Industry Research Assistance Council (BIRAC) for their invaluable financial support and provision of laboratory facilities. This underscores the collaborative efforts in advancing healthcare innovation. Special gratitude is offered to the Department of Biochemistry at AIIMS Bhopal for generously providing

essential facilities to conduct the wound healing assay, cytotoxicity tests, and scratch tests. Their support has played a pivotal role in successfully executing our research.

## CONTRIBUTION OF AUTHORS

**Research concept-** Divyanshi Makode, Vishvam Jawade and Mahi Verma

**Research design-** Dr. Nidhi Tripathi

**Supervision-** Prof. Jagat Rakesh Kanwar and Prof. Rashmi Chowdhary

**Materials-** Ashish Kumar Yadav and Vrushali D Patil

**Data collection-** Ashish Kumar Yadav

**Data analysis and Interpretation-** Ashish Kumar Yadav and Vrushali D Patil

**Literature search-** Divyanshi Makode, Vishvam Jawade and Mahi Verma

**Writing article-** Dr. Nidhi Tripathi

**Critical review-** Prof. Jagat Rakesh Kanwar and Prof. Rashmi Chowdhary

**Article editing-** Dr. Nidhi Tripathi

**Final approval-** Dr. Anjali Choudhary and Dr. Preeti Chandurkar

## REFERENCES

- [1] Elzayat EM, Auda, SH, Alanazi, FK, Al-Agamy, MH. Evaluation of wound healing activity of henna, pomegranate and myrrh herbal ointment blend. *Saudi Pharm J.*, 2018; 26(5): 733-38.
- [2] Gonzalez ACDO, Andrade ZDA, Costa TF, Medrado ARAP Wound healing—A literature review. *Anais Brasileiros de Dermatologia*, 2016; 91: 614.
- [3] Xue M, Jackson, CJ. Extracellular matrix reorganization during wound healing and its impact on abnormal scarring. *Adv Wound Care*, 2015; 4: 119–36.
- [4] Ghavide ND, Naderi MS, Tabaie SM. Role of proteins and effective factors in wound healing. *J Lasers Med Sci.*, 2018; 2: 35–48.
- [5] Mulkalwar S, Behera L, Golande P, Manjare R, Patil H. Evaluation of wound healing activity of topical phenytoin in an excision wound model in rats. *Int J Basic Clin Pharmacol.*, 2015; 4: 139–43.
- [6] Shrimanker M, Patel N, Modi H, Dave R. A review: screening models for wound healing activity in animals. *Am J Pharm Tech Res.*, 2013; 3: 237–51.

- [7] Velnar T, Bailey T, Smrkolj V. The wound healing process: an overview of the cellular and molecular mechanisms. *J Int Med Res.*, 2009; 37: 1528–42.
- [8] Gosain A, Dipietro LA. Aging and wound healing. *World J Surg.*, 2004; 28: 321–26.
- [9] Farahpour MR, Mirzakhani N, Doostmohammadi J, Ebrahimzadeh M. Hydroethanolic *Pistacia atlantica* hulls extract improved wound healing process; evidence for mast cells infiltration, angiogenesis and RNA stability. *Int J Surg.*, 2015; 17: 88–98.
- [10] Farahpour MR, Sedaghat S. Effect of *Malva Sylvestris* hydroethanolic leaf extract on the healing of full-thickness, excisional skin wounds in the rat. *Veterinary Clin Pathol.*, 2015; 1: 73–81.
- [11] Segeritz CP, Vallier L. Cell Culture: Growing Cells as Model Systems *In Vitro*. *Basic Sci Methods Clin Res.*, 2017; pp. 151–72.
- [12] Alankar S, Arun KM, Abid M, Aftab et al. Extracts of *Tridax procumbens* Linn leaves causes wound healing in diabetic and non-diabetic laboratory animals. *Wound Med.*, 2020; 29: 100185.
- [13] Ganesan P, Pradeepa, P. Development and Characterization of nanofibrous mat from PVA/*Tridax Procumbens* (TP) leaves extracts. *Wound Med.*, 2017; 19: 15-22.
- [14] Trupti S, Prachi B, Prajakta D, Ravindra A. Fabrication of Herbal Hemostat films loaded with medicinal *Tridax procumbens* extracts. *Fibers Polymers*, 2021; 22(8): 2135-44.
- [15] Jude CI. An aqueous extract of the leaves of *Tridax procumbens* Linn (Asteraceae) protected against carbon tetrachloride-induced liver injury in Wistar rats. *Pacific J Sci Technol.*, 2012; 13(1): 519-27.
- [16] Radstake WE, Gautam K, Van RC, et al. Comparison of in vitro scratch wound assay experimental procedures. *Biochem Biophys Reports*, 2023; 33: 101423.
- [17] Paralkar S, Kamalapurkar KA, Koli L. Formulation and Evaluation of *Tridax Procumbens* Herbal Gel. *Int J Res Pharma Biomed Sci.*, 2020; 8(2): 110-13.
- [18] Roy IM, Karuppur T, et al. Evaluation of Wound Healing Property of the Ethanolic Extract of *Glycyrrhiza Glabra* on Vero Cell Lines Using *In Vitro* Scratch Assay Test. *J Pharm Bioallied Sci.*, 2023; 15(S1): S630-S35.
- [19] Benito D. Johnson and Appalaraju Gorle. Evaluation of Ethanolic Extract of *Tridax procumbens* Linn on Wound Healing in Rats. *Res J Pharm Tech.*, 2012; 5(2): 239-42.
- [20] Muthukumar B, Nandini MS, Elumalai P, et al. Enhancement of cell migration and wound healing by nano-herb ointment formulated with biosurfactant, silver nanoparticles and *Tridax procumbens*. *Front Microbiol.* 2023; 2.
- [21] Nimbekar TP, Meshram AS, Farooqui MS. Formulation and evaluation of ointment containing natural wound healing activity of *Tridax procumbens*. *J Emer Tech Inno Res.*, 2019; 6(5): 562.
- [22] Talekar, YP, Apte KG, Paygude SV, et al. Studies on wound healing potential of polyherbal formulation using *in vitro* and *in vivo* assays. *J Ayurveda Integr Med.*, 2017; (8): 73-81.
- [23] Venkatachalam H, Palaniswamy R. Evaluation of *Tridax Procumbens* Leaf Extract Loaded PVA Film for Wound Healing Application. *Int J Pharm Sci Res.*, 2023; 14(9): 4474-80.

**Open Access Policy:**

Authors/Contributors are responsible for originality, contents, correct references, and ethical issues. SSR-IJLS publishes all articles under Creative Commons Attribution- Non-Commercial 4.0 International License (CC BY-NC). <https://creativecommons.org/licenses/by-nc/4.0/legalcode>

